

Classical Gravitational Interactions and Gravitational Lorentz Force

Ning Wu *

Institute of High Energy Physics, P.O.Box 918-1, Beijing 100039, P.R.China

February 7, 2008

PACS Numbers: 11.15.-q, 04.60.-m.

Keywords: gravitational Lorentz force, gravity, gauge field, quantum gravity.

Abstract

In quantum gauge theory of gravity, the gravitational field is represented by gravitational gauge field. The field strength of gravitational gauge field has both gravitoelectric component and gravitomagnetic component. In classical level, gauge theory of gravity gives out classical Newtonian gravitational interactions in a relativistic form. Besides, it gives out gravitational Lorentz force which is the gravitational force on a moving object in gravitomagnetic field. The direction of gravitational Lorentz force does not along that of classical gravitational Newtonian force. Effects of gravitational Lorentz force should be detectable, and these effects can be used to discriminate gravitomagnetic field from ordinary electromagnetic magnetic field.

*email address: wuning@mail.ihep.ac.cn

1 Introduction

In classical Newton's theory of gravity, gravity obeys the inverse square law[1]. That is, the gravitational force between two mass point m_1 and m_2 is

$$f = \frac{G_N m_1 m_2}{r^2}, \quad (1.1)$$

where G_N is the newtonian gravitational constant and r is the distance between two mass point. Gravitational force is along the line which connects the mass point and is always attractive. Obvious, the above expression is not invariant or covariant under Lorentz transformation.

In Einstein's general theory of relativity[2, 3], gravity is treated as space-time geometry, and gravity is just an effect of the curved space-time. General relativity is a geometric theory of gravity, and in the original expressions, the concepts of the space-time metric, affine connection, curvature tensor, geodesic curve, $\cdots$ are used, and the concept of the traditional "gravitational force" is not used. Only in post-Newtonian approximation can we clearly see the correspondent of traditional gravitational force and its influences[4, 5].

Quantum gauge theory of gravity is proposed in the framework of the quantum field theory[6, 7, 8, 9, 10], where gravity is treated as a kind of physical interactions and space-time is always kept flat. This treatment satisfies the fundamental spirit of quantum field theory. Quantum gauge theory of gravity is a perturbatively renormalizable quantum theory. Quantum gauge theory can be used to solve some problems related to quantum behavior of gravitational interactions, such as unifications of fundamental interactions can be easily accomplished in it[12, 13, 14], it can be used to explain the possible origin of dark matter and dark energy[15, 16], it can be used to explain Podkletnov effects[17] and COW experiments.

In gauge theory of gravity, gravitoelectromagnetic field is naturally defined as components of field strength of gravitational gauge field. It is known that gravitoelectromagnetism was studied for more than 130 years. The close analogy between Newton's gravitation law and Coulomb's law of electricity led to the birth of the concept of gravitomagnetism in the nineteenth century[18, 19, 20, 21]. Later, in the framework of General Relativity, gravitomagnetism was extensively explored[22, 23, 24] and recently some experiments are designed to test gravitomagnetism effects. Some recently reviews on gravitomagnetism can be found in literatures [25, 26, 27]. In quantum gauge theory of gravity, gravitoelectromagnetism is defined in a more general way, and the gravitoelectromagnetism discussed in the literatures [22, 23, 24, 25, 26, 27] is only the special case of it.

In this paper, we will discuss some classical effects of gravitational interactions in the framework of gauge theory of gravity, including gravitational interactions of two mass point and gravitational Lorentz force.

2 Pure Gravitational Gauge Field

First, for the sake of integrity, we give a simple introduction to gravitational gauge theory and introduce some notations which is used in this paper. Details on quantum gauge theory of gravity can be found in literatures [6, 7, 8, 9, 10]. In gauge theory of gravity, the most fundamental quantity is gravitational gauge field $C_\mu(x)$, which is the gauge potential corresponding to gravitational gauge symmetry. Gauge field $C_\mu(x)$ is a vector in the corresponding Lie algebra, which is called the gravitational Lie algebra. So $C_\mu(x)$ can expanded as

$$C_\mu(x) = C_\mu^\alpha(x) \hat{P}_\alpha, \quad (\mu, \alpha = 0, 1, 2, 3) \quad (2.1)$$

where $C_\mu^\alpha(x)$ is the component field and $\hat{P}_\alpha = -i\frac{\partial}{\partial x^\alpha}$ is the generator of the global gravitational gauge group. The gravitational gauge covariant derivative is defined by

$$D_\mu = \partial_\mu - igC_\mu(x) = G_\mu^\alpha \partial_\alpha, \quad (2.2)$$

where g is the gravitational coupling constant and G is given by

$$G = (G_\mu^\alpha) = (\delta_\mu^\alpha - gC_\mu^\alpha) = (I - gC)_\mu^\alpha. \quad (2.3)$$

Matrix G is an important quantity in gauge theory of gravity. Its inverse matrix is denoted as G^{-1}

$$G^{-1} = \frac{1}{I - gC} = (G_\alpha^{-1\mu}). \quad (2.4)$$

Using matrix G and G^{-1} , we can define two important quantities

$$g^{\alpha\beta} = \eta^{\mu\nu} G_\mu^\alpha G_\nu^\beta, \quad (2.5)$$

$$g_{\alpha\beta} = \eta_{\mu\nu} G_\alpha^{-1\mu} G_\beta^{-1\nu}. \quad (2.6)$$

The field strength of gravitational gauge field is defined by

$$F_{\mu\nu} \triangleq \frac{1}{-ig} [D_\mu, D_\nu]. \quad (2.7)$$

Its explicit expression is

$$F_{\mu\nu}(x) = \partial_\mu C_\nu(x) - \partial_\nu C_\mu(x) - igC_\mu(x)C_\nu(x) + igC_\nu(x)C_\mu(x). \quad (2.8)$$

$F_{\mu\nu}$ is also a vector in gravitational Lie algebra and can be expanded as,

$$F_{\mu\nu}(x) = F_{\mu\nu}^\alpha(x) \cdot \hat{P}_\alpha, \quad (2.9)$$

where

$$F_{\mu\nu}^\alpha = \partial_\mu C_\nu^\alpha - \partial_\nu C_\mu^\alpha - gC_\mu^\beta \partial_\beta C_\nu^\alpha + gC_\nu^\beta \partial_\beta C_\mu^\alpha. \quad (2.10)$$

3 Gravitoelectromagnetic Field

$F_{\mu\nu}^\alpha$ is the component field strength of gravitational gauge field. Define

$$F_{ij}^\alpha = -\varepsilon_{ijk} B_k^\alpha, \quad F_{0i}^\alpha = E_i^\alpha. \quad (3.1)$$

Then field strength of gravitational gauge field can be expressed as

$$F^\alpha = \begin{pmatrix} 0 & E_1^\alpha & E_2^\alpha & E_3^\alpha \\ -E_1^\alpha & 0 & -B_3^\alpha & B_2^\alpha \\ -E_2^\alpha & B_3^\alpha & 0 & -B_1^\alpha \\ -E_3^\alpha & -B_2^\alpha & B_1^\alpha & 0 \end{pmatrix}. \quad (3.2)$$

This form is quite similar to that of field strength in electrodynamics, but with an extra group index α . The component E_i^α of field strength is called gravitoelectric field, and B_i^α is called gravitomagnetic field. Traditional Newtonian gravity is transmitted by E_i^0 . The $\alpha = 0$ components B_i^0 and E_i^0 correspond to the gravitoelectromagnetic field defined in literature [22, 23, 24, 25, 26, 27].

In most cases, gravitational field is weak and its self interactions can be neglected. So, eq.(2.10) can be simplified to

$$F_{\mu\nu}^\alpha = \partial_\mu C_\nu^\alpha - \partial_\nu C_\mu^\alpha. \quad (3.3)$$

The gravitomagnetic and gravitoelectric field have much familiar forms

$$B_i^\alpha = \partial_k C_j^\alpha - \partial_j C_k^\alpha, \quad (3.4)$$

$$E_i^\alpha = \partial_t C_i^\alpha - \partial_i C_0^\alpha \quad (3.5)$$

From definitions eq.(3.1), we can prove that

$$\nabla \cdot \vec{B}^\alpha = 0, \quad (3.6)$$

$$\frac{\partial}{\partial t} \vec{B}^\alpha + \nabla \times \vec{E}^\alpha = 0. \quad (3.7)$$

From eq.(3.6), we know that gravitomagnetic field is source free. It is found that gravitomagnetic field is generated by moving objects, and transmit gravitomagnetic interactions between two rotating objects.

4 Gravitational Lorentz Force

A particle which moves in a gravitomagnetic field will feel a force that is perpendicular to its motion. In electrodynamics, this force is usually called Lorentz force. As an example, we discuss gravitational interactions between gravitational field and Dirac field. The lagrangian for gravitational gauge interactions of Dirac field is[28, 11]

$$\mathcal{L}_0 = -\bar{\psi}(\gamma^\mu D_\mu + m)\psi - \frac{1}{4}C_{\alpha\beta}^{\mu\nu\rho\sigma}F_{\mu\nu}^\alpha F_{\rho\sigma}^\beta, \quad (4.1)$$

where

$$C_{\alpha\beta}^{\mu\nu\rho\sigma} = \frac{1}{4}\eta^{\mu\rho}\eta^{\nu\sigma}g_{\alpha\beta} + \frac{1}{2}\eta^{\mu\rho}G_\beta^{-1\nu}G_\alpha^{-1\sigma} - \eta^{\mu\rho}G_\alpha^{-1\nu}G_\beta^{-1\sigma}. \quad (4.2)$$

So, the interaction Lagrangian is

$$\mathcal{L}_I = g\bar{\psi}\gamma^\mu\partial_\alpha\psi C_\mu^\alpha. \quad (4.3)$$

For Dirac field, the gravitational energy-momentum of Dirac field is

$$T_{g\alpha}^\mu = \bar{\psi}\gamma^\mu\partial_\alpha\psi. \quad (4.4)$$

Substitute eq.(4.4) into eq.(4.3), we get

$$\mathcal{L}_I = gT_{g\alpha}^\mu C_\mu^\alpha. \quad (4.5)$$

The interaction Hamiltonian density $\mathcal{H}_I$ is

$$\mathcal{H}_I = -\mathcal{L}_I = -gT_{g\alpha}^\mu(y, \vec{x})C_\mu^\alpha(y). \quad (4.6)$$

Suppose that the moving particle is a mass point at point $\vec{x}$, in this case

$$T_{g\alpha}^\mu(y, \vec{x}) = T_{g\alpha}^\mu\delta(\vec{y} - \vec{x}), \quad (4.7)$$

where $T_{g\alpha}^\mu$ is independent of space coordinates. Then, the interaction Hamiltonian H_I is

$$H_I = \int d^3 \vec{y} \mathcal{H}_I(y) = -g \int d^3 \vec{y} T_{g\alpha}^\mu(y, \vec{x}) C_\mu^\alpha(y). \quad (4.8)$$

The gravitational force that acts on the mass point is

$$f_i = g \int d^3 y T_{g\alpha}^\mu(y, \vec{x}) F_{i\mu}^\alpha + g \int d^3 y T_{g\alpha}^\mu(y, \vec{x}) \frac{\partial}{\partial y^\mu} C_i^\alpha. \quad (4.9)$$

For quasi-static system, if we omit higher order contributions, the second term in the above relation vanish. For a mass point, using the technique of Lorentz covariance analysis, we can prove that

$$P_{g\alpha} U^\mu = \gamma T_{g\alpha}^\mu, \quad (4.10)$$

where U^μ is velocity, γ is the rapidity, and $P_{g\alpha}$ is the gravitational energy-momentum. According to eq.(4.7), $P_{g\alpha}$ is given by

$$P_{g\alpha} = \int d^3 \vec{y} T_{g\alpha}^0(y) = T_{g\alpha}^0. \quad (4.11)$$

Using all these relations and eq.(3.1), we get

$$\vec{f} = -g P_{g\alpha} \vec{E}^\alpha - g P_{g\alpha} \vec{v} \times \vec{B}^\alpha. \quad (4.12)$$

For quasi-static system, the dominant contribution of the above relation is

$$\vec{f} = g M \vec{E}^0 + g M \vec{v} \times \vec{B}^0, \quad (4.13)$$

where $\vec{v} = \vec{U} / \gamma$ is the velocity of the mass point. The first term of eq.(4.13) is the classical Newton's gravitational interactions. The second term of eq.(4.13) is the Lorentz force. The direction of this force is perpendicular to the direction of the motion of the mass point. When the mass point is at rest or is moving along the direction of the gravitomagnetic field, this force vanishes. Lorentz force should have some influences for cosmology, for the rotation of galaxy will generate gravitomagnetic field and this gravitomagnetic field will affect the motion of stars and affect the large scale structure of galaxy.

5 Repulsive Component of Gravitational Interactions

The classical gravitational interactions are always attractive, but sometimes, there are repulsive components in gravity. The gravitational force is given by eq.(4.12).

The first term corresponds to classical gravitational force. It is

$$\begin{aligned}\vec{f} &= -gP_{g\alpha} \vec{E}^\alpha \\ &= g(M_1 \vec{E}^0 - P_{gj} \vec{E}^j),\end{aligned}\tag{5.1}$$

where M_1 is the gravitational mass of the mass point which is moving in gravitational field. Suppose that the gravitational field is generated by another mass point whose gravitational energy-momentum is Q_g^α and its gravitational mass is M_2 . For quasi-static gravitational field, we can get

$$\vec{E}^\alpha = -\frac{g}{4\pi r^3} Q_g^\alpha \vec{r}\tag{5.2}$$

Substitute eq.(5.2) into eq.(5.1), we get

$$\vec{f} = \frac{g^2}{4\pi r^3} \vec{r} (-E_{1g}E_{2g} + \vec{P}_g \cdot \vec{Q}_g),\tag{5.3}$$

where E_{1g} and E_{2g} are gravitational energy of two mass point. From eq.(5.3), we can see that, if $\vec{P}_g \cdot \vec{Q}_g$ is positive, the corresponding component gravitational force between two momenta is repulsive. Because $E_{1g}E_{2g} \geq \vec{P}_g \cdot \vec{Q}_g$, the total gravitational force between two mass point is always attractive.

Suppose that a Lorentz transformation along the direction which is perpendicular to the direction of the gravitational force $\vec{f}$, the left hand side of eq.(5.3) is invariant. Under this transformation, the gravitational energy of both mass point will be changed, so $E_{1g}E_{2g}$ will be changed, but the sum $-E_{1g}E_{2g} + \vec{P}_g \cdot \vec{Q}_g$ keep invariant. Therefore, the appearance of the component gravitational force between two momenta satisfies the requirement of Lorentz symmetry.

6 Discussions

Eq.(4.12) is obtained from the interaction Lagrangian, it is deduced without concerning dynamics of gravitational gauge field. It is known that the selection of the Lagrangian of pure gravitational gauge field is not unique, but this ambiguity does not affect Eq.(4.12). Different selection of the Lagrangian of pure gravitational gauge field only gives out different $\vec{E}^\alpha$ and $\vec{B}^\alpha$, it does not affect classical Newtonian gravitational interactions and gravitational Lorentz force.

In classical Newton's theory of gravity, gravitational force between two objects is always along the line which connects the center of mass of the two objects. Because of the gravitational Lorentz force, there exists components force which is perpendicular to the line. In most cases, this orthogonal component is much weaker than the traditional Newtonian gravity.

Gravitational Lorentz force is independent of electric charge, so it is different from the electric Lorentz force. Only charged particle can feel electric Lorentz force, but all particles can feel gravitational Lorentz force. Using this property, we can discriminate gravitational Lorentz force from electric Lorentz force, and discriminate gravitomagnetic field from the electromagnetic magnetic field. Effects of gravitational Lorentz force should be observable and can be observed by astrophysical observations. It is known that there are some experiments designed to test gravitomagnetic effects.

References

- [1] Isaac Newton, *Mathematical Principles of Natural Philosophy*, (Cambridge University Press, 1934).
- [2] Albert Einstein, *Annalen der Phys.*, **49** (1916) 769 .
- [3] Albert Einstein, *Zeits. Math. und Phys.* **62** (1913) 225.
- [4] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*, (John Wiley, New York, 1972).
- [5] N. Straumann, *General Relativity and Relativistic Astrophysics*, (Springer-Verlag, Berlin Heidelberg New York Tokyo, 1984).
- [6] Ning Wu, "Gauge Theory of Gravity", hep-th/0109145
- [7] Ning Wu, *Commun. Theor. Phys. (Beijing, China)* **38** (2002): 151-156.
- [8] Ning Wu, "Quantum Gauge Theory of Gravity", hep-th/0112062
- [9] Ning Wu, "Quantum Gauge Theory of Gravity", talk given at Meeting of the Division of Particles and Fields of American Physical Society at the College of William & Mary(DPF2002), May 24-28, 2002, Williamsburg, Virgia, USA; hep-th/0207254; Transparency can be obtained from: http://dpf2002.velopers.net/talks_pdf/33talk.pdf

- [10] Ning Wu, "Renormalizable Quantum Gauge General Relativity" gr-qc/0309041.
- [11] Ning Wu, Commun. Theor. Phys. (Beijing, China) **42** (2004): 543-552.
- [12] Ning Wu, Commun. Theor. Phys. (Beijing, China) **38** (2002): 322-326.
- [13] Ning Wu, Commun. Theor. Phys. (Beijing, China) **38** (2002): 455-460.
- [14] Ning Wu, Commun. Theor. Phys. (Beijing, China) **39** (2003): 561-568.
- [15] Ning Wu, Commun. Theor. Phys. (Beijing, China) **36**(2001) 169-172.
- [16] Ning Wu, Commun. Theor. Phys. (Beijing, China) **39** (2003): 671-674.
- [17] Ning Wu, Commun. Theor. Phys. (Beijing, China) **41**(2004) 567-572; hep-th/0307225.
- [18] Maxwell J.C., Phil. Trans., **155**, (1865) 492.
- [19] G. Holzmüller, Z. Math. Phys., **15**, (1870) 69.
- [20] F. Tisserand, Compte Rendu hebdomadaire des seances de L'Academie des sciences, **75** (1872) 760; **110** (1890) 313.
- [21] O. Heaviside, Electromagnetic Theory, The Electrician Printing and Publishing Co., London, 1894, Vol. I, Appendix B; O. Heaviside, The Electrician, **31** (1893) 281-282 and 359.
- [22] A. Einstein, Phys. Z., **14**, (1913) 1261.
- [23] H. Thirring, Phys. Z., **19**, (1918) 204; **19**, (1918) 33; **22**, (1921) 29.
- [24] J. Lense and H. Thirring, Phys. Z., **19**, (1918) 156; B. Mashhoon, F.W. Hehl and D.S. Thesis, Gen. Rel. Grav., **16**, (1984) 711.
- [25] M.L. Ruggiero and A. Tartaglia, gr-qc/0207065.
- [26] B. Mashhoon, gr-qc/0311030.
- [27] G. Schafer, gr-qc/0407116.
- [28] Ning Wu, Commun. Theor. Phys. (Beijing, China) **41**(2004) 381-384;